

Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

SONATA®

Control of Leaf spot of Sugar Beet when tank mixed with another registered fungicide.

EPA Reg. No. 264-1153

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF MI, MN, and ND

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

Sonata is recommended to control Leaf Spot of Sugar Beet when tank mixed with another registered fungicide.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. For use only as described on this label. Not for isolation or deformulation. Do not culture.

SONATA may be tank-mixed with other registered pesticides to enhance plant disease control. This product cannot be mixed with any product with a prohibition against such mixing. When tank-mixing SONATA with other registered pesticides, always read and follow all use directions, restrictions, and precautions of both SONATA and the tank-mix partner(s). Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions. Do not exceed label dosage rates.

Crops	Disease	Rate qt/acre	Application Instructions
Sugar Beets Including: Crop grown for seed production	Leaf Spot Cercospora beticola	1-4	Begin applications when environmental conditions are conducive to disease development. Continue applications on 7- to 14-day intervals or as needed. Use the stated higher rates and shorter application intervals under heavy disease pressure. The lowest rate of 1 qt/acre is only for use in tank mixes with other registered fungicides

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577. For <u>PRODUCT USE</u> Information Call 1-866-99BAYER (1-866-992-2937).

> As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).